



# Essai Clinique

Généré le 04 mai 2024 à partir de

Titre	A Randomized Phase 3 Study of the Combination of Pembrolizumab Plus Epacadostat Alone or With Platinum-based Chemotherapy Versus Pembrolizumab Plus Platinum-based Chemotherapy Plus Placebo as First-Line Treatment in Patients With Metastatic Non-Small Cell Lung Cancer
Protocole ID	MK-3475-715/KEYNOTE-715-05
ClinicalTrials.gov ID	<a href="https://clinicaltrials.gov/ct2/show/study/NCT03322566">NCT03322566</a>
Type(s) de cancer	Poumon non à petites cellules
Phase	Phase III
Stade	Métastatique
Type étude	Traitement
Médicament	Epacadostat, Pembrolizumab, Chimio
Institution	CIUSSS DE L'OUEST-DE-L'ILE-DE-MONTREAL CENTRE HOSPITALIER DE ST. MARY 3830 av. Lacombe, Montréal, QC, H3T 1M5
Ville	Montreal
Investigateur principal	Dr Adrian Langleben
Coordonnateur	Dora Bartulovic 514-345-3511 poste 3378
Statut	Fermé
But étude	The purpose of this study is to evaluate the efficacy and safety of pembrolizumab plus epacadostat alone or with platinum-based chemotherapy versus pembrolizumab plus platinum-based chemotherapy plus placebo as first-line therapy in participants with metastatic non-small cell lung cancer (NSCLC).
Critères d'éligibilité	<ul style="list-style-type: none"><li>• Histologically or cytologically confirmed diagnosis of stage IV NSCLC without epidermal growth factor receptor (EGFR)-sensitizing mutation, ROS1 and/or anaplastic lymphoma kinase (ALK) translocation</li><li>• Measurable disease based on RECIST 1.1</li><li>• Life expectancy of at least 3 months.</li><li>• Eastern Cooperative Oncology Group (ECOG) performance status of 0 or 1.</li><li>• Adequate organ function per protocol-defined criteria.</li><li>• Provide tumor tissue sample.</li></ul>
Critères d'exclusion	<ul style="list-style-type: none"><li>• Known untreated central nervous system metastases and/or carcinomatous meningitis</li><li>• History of (non-infectious) pneumonitis that required systemic steroids or current pneumonitis/interstitial lung disease.</li><li>• Symptomatic ascites or pleural effusion.</li><li>• Known history of an additional malignancy, except if the participant has undergone potentially curative therapy with no evidence of that disease recurrence for 5 years since initiation of that therapy.</li><li>• Active autoimmune disease that has required systemic treatment in past 2 years.</li><li>• Has had an allogeneic tissue/solid organ transplant.</li><li>• Has a known history of human immunodeficiency virus (HIV) infection. HIV testing is not required unless mandated by the local health authority.</li><li>• Has known history of or is positive for active Hepatitis B (HBsAg reactive) or has active Hepatitis C (HCV RNA). Note: Testing must be performed to determine eligibility.</li><li>• History or presence of an abnormal electrocardiogram (ECG) that, in the Investigator's opinion,</li></ul>

is clinically meaningful.

- Use of protocol-defined prior/concomitant therapy.