



# Essai Clinique

Généré le 28 avr. 2024 à partir de

Titre	Étude de phase 2, de l'INCMGA00012 chez des participants atteints d'un carcinome à cellules de Merkel métastatique
Protocole ID	INCMGA 0012-201
ClinicalTrials.gov ID	<a href="#">NCT03599713</a>
Type(s) de cancer	Peau
Phase	Phase II
Stade	Métastatique
Type étude	Traitement
Médicament	INCMGA00012
Institution	CIUSSS DU CENTRE-OUEST-DE-L'ILE-DE-MONTREAL HOPITAL GENERAL JUIF SIR MORTIMER B.DAVIS 3755 rue de la Côte Ste. Catherine, Montréal, QC, H3T 1E2
Ville	Montréal
Investigateur principal	Dr Wilson Miller
Coordonnateur	Ralph Waked 514-340-8222 poste 26823
Statut	Fermé
But étude	Le but de cette étude est d'évaluer l'activité clinique et l'innocuité de l'INCMGA00012 chez les participants atteints d'un carcinome métastatique à cellules de Merkel.
Critères d'éligibilité	<ul style="list-style-type: none"><li>• Signed informed consent.</li><li>• Diagnosis of MCC with distant metastatic disease as a component of tumor burden and no more than 3 prior systemic treatments, inclusive of systemic adjuvant therapy.</li><li>• Eastern Cooperative Oncology Group performance status of 0 to 1.</li><li>• Measurable disease according to RECIST v1.1.</li><li>• Availability of tumor tissue (fresh or archival) for central pathology review.</li><li>• Willingness to avoid pregnancy or fathering children based on protocol-defined criteria.</li></ul>
Critères d'exclusion	<ul style="list-style-type: none"><li>• Prior programmed cell death protein 1 (PD-1) or programmed cell death ligand protein 1 (PD-L1)-directed therapy.</li><li>• Treatment with anticancer drugs or participation in another interventional clinical study within 21 days before the first administration of study drug.</li><li>• Has not recovered to ≤ Grade 1 or baseline from toxic effects of prior therapy (with the exceptions for anemia not requiring transfusion support and any grade of alopecia) and/or complications from prior surgical intervention within 7 days before starting study treatment.</li><li>• Radiation therapy administered within 2 weeks of first dose of study treatment or radiation therapy in the thoracic region that is &gt; 30 Gy within 6 months of the first dose of study treatment.</li><li>• Known central nervous system (CNS) metastases and/or carcinomatous meningitis.</li><li>• History of second malignancy within 3 years (with exceptions).</li><li>• Laboratory values outside the protocol-defined range at screening.</li><li>• Clinically significant pulmonary, cardiac, gastrointestinal or autoimmune disorders.</li><li>• Active bacterial, fungal, or viral infections, including hepatitis A, B, and C.</li><li>• Receipt of a live vaccine within 90 days of planned start of study therapy.</li><li>• Current use of protocol-defined prohibited medication.</li><li>• Known hypersensitivity to another monoclonal antibody that cannot be controlled with standard</li></ul>

measures (eg, antihistamines and corticosteroids).

- Inability or unlikely, in the opinion of the investigator, to comply with the Protocol requirements.
- Participant who is pregnant or breastfeeding.