



# Clinical Trial

Generated on 25 Apr 2025 from the

Title	A Phase 1/2 Study of BMS-986340 as Monotherapy and in Combination With Nivolumab in Participants With Advanced Solid Tumors
Protocole ID	CA052-002
ClinicalTrials.gov ID	<a href="#">NCT04895709</a>
Cancer Type(s)	Tumeurs solides
Phase	Phase I-II
Stage	Maladie avancée ou métastatique
Study Type	Clinique
Drug	BMS-986340 en monothérapie ou en association avec le nivolumab
Institution	CENTRE HOSPITALIER DE L'UNIVERSITE DE MONTREAL
City	
Principal Investigator	
Coordinator	
Status	Actif en recrutement
Eligibility Criteria	<ul style="list-style-type: none"><li>• Fresh pre-treatment and on-treatment tumor biopsy must be provided for biomarker analysis</li><li>• Measurable disease per Response Evaluation Criteria in Solid Tumors (RECIST) v1.1 and at least 1 lesion accessible for biopsy</li><li>• Eastern Cooperative Oncology Group Performance Status of 0 or 1</li><li>• Radiographically documented progressive disease on or after the most recent therapy</li><li>• Received standard-of-care therapies, including an available programmed death (ligand)-1 inhibitor known to be effective in the tumor type for which they are being evaluated</li><li>• Parts 1A, 1B, and 2A: Advanced or metastatic non-small cell lung cancer, squamous cell carcinoma of head and neck, microsatellite stable colorectal cancer, gastric/ gastroesophageal junction adenocarcinoma, or cervical cancer, and have received, be refractory to, not be a candidate for, or be intolerant of existing therapies known to provide clinical benefit for the condition of the participant</li></ul>
Exclusion Criteria	<ul style="list-style-type: none"><li>• Women who are pregnant or breastfeeding</li><li>• Primary central nervous system (CNS) malignancy</li><li>• Untreated CNS metastases</li><li>• Leptomeningeal metastases</li><li>• Concurrent malignancy requiring treatment or history of prior malignancy active within 2 years prior to the first dose of study treatment</li><li>• Active, known, or suspected autoimmune disease</li><li>• Condition requiring systemic treatment with either corticosteroids within 14 days or other immunosuppressive medications within 30 days of the first dose of study treatment</li><li>• Prior organ or tissue allograft</li><li>• Uncontrolled or significant cardiovascular disease</li><li>• Major surgery within 4 weeks of study drug administration</li><li>• History of or with active interstitial lung disease or pulmonary fibrosis</li></ul> <p>Other protocol-defined inclusion/exclusion criteria apply</p>