



# Essai Clinique

Généré le 12 mai 2025 à partir de

Titre	A Phase 2 Study of VS-6766 (Dual RAF/MEK Inhibitor) Alone and In Combination With Defactinib (FAK Inhibitor) in Recurrent Low-Grade Serous Ovarian Cancer (LGSOC)
Protocole ID	RAMP-201 (ENGOT-ov60)
ClinicalTrials.gov ID	<a href="https://clinicaltrials.gov/ct2/show/study/NCT04625270">NCT04625270</a>
Type(s) de cancer	Ovaire
Phase	Phase II
Type étude	Clinique
Médicament	VS-6766 seul ou en association avec defactinib
Institution	CENTRE HOSPITALIER DE L'UNIVERSITE DE MONTREAL
Ville	
Investigateur principal	Dre Diane Provencher
Coordonnateur	Bonny Choy 514-890-8000 poste 24672
Statut	Fermé
But étude	This study will assess the safety and efficacy of VS-6766 monotherapy and in combination with Defactinib in subjects with recurrent Low-Grade Serous Ovarian Cancer (LGSOC)
Critères d'éligibilité	<ul style="list-style-type: none"><li>• Histologically proven LGSOC (ovarian, peritoneal)</li><li>• In Part A KRAS mutation, KRAS wt</li><li>• Progression or recurrence of LGSOC after at least one prior systemic therapy for metastatic disease.</li><li>• Measurable disease according to RECIST 1.1</li><li>• An Eastern Cooperative Group (ECOG) performance status <math>\leq 1</math>.</li><li>• Adequate organ function</li><li>• Adequate recovery from toxicities related to prior treatments</li><li>• Agreement to use highly effective method of contraceptive</li></ul>
Critères d'exclusion	<ul style="list-style-type: none"><li>• Systemic anti-cancer therapy within 4 weeks of the first dose of study therapy</li><li>• Co-existing high-grade ovarian cancer or another histology</li><li>• History of prior malignancy with recurrence <math>&lt;3</math> years from the time of enrollment</li><li>• Major surgery within 4 weeks</li><li>• Symptomatic brain metastases requiring steroids or other interventions</li><li>• Known SARS-Cov2 infection (clinical symptoms) <math>\leq 28</math> days prior to first dose of study therapy</li><li>• For subjects with prior MEK exposure, Grade 4 toxicity deemed related to the MEK inhibitor</li><li>• Active skin disorder that has required systemic therapy within the past year</li><li>• History of rhabdomyolysis</li><li>• Concurrent ocular disorders</li><li>• Concurrent heart disease or severe obstructive pulmonary disease</li><li>• Subjects with the inability to swallow oral medications</li></ul>