


Titre	A phase 3, randomized, open-label, multicenter trial of ARV-471 (PF-07850327) vs fulvestrant in participants with estrogen receptor-positive, her2-negative advanced breast cancer whose disease progressed after prior endocrine based treatment for advanced disease
Protocole ID	VERITAC-2
ClinicalTrials.gov ID	<a href="https://clinicaltrials.gov/ct2/show/study/NCT05654623">NCT05654623</a>
Type(s) de cancer	Sein
Phase	Phase III
Type étude	Clinique
Médicament	ARV-471 (PF-07850327) versus fulvestrant
Institution	CISSE DES LAURENTIDES  HOPITAL DE SAINT-JEROME 290 Rue de Montigny, Saint-Jérôme, QC, J7Z 5T3
Ville	
Investigateur principal	Dr Ghislain Cournoyer
Coordonnateur	Yanick Sardin Laframboise 450-431-1020 poste 23429
Statut	Actif en recrutement
But étude	A study to learn about a new medicine called ARV-471 (PF-07850327) in people who have advanced metastatic breast cancer.
Critères d'éligibilité	<ul style="list-style-type: none"><li>• Adult participants with loco-regional recurrent or metastatic breast disease not amenable to surgical resection or radiation therapy</li><li>• Confirmed diagnosis of ER+/HER2- breast cancer</li><li>• Prior therapies for locoregional recurrent or metastatic disease must fulfill all the following criteria:<ul style="list-style-type: none"><li>• One line of CDK4/6 inhibitor therapy in combination with endocrine therapy. Only one line of CDK4/6 inhibitor is allowed in any setting.</li><li>• ≤ 1 endocrine therapy in addition to CDK4/6 inhibitor with ET</li><li>• Most recent endocrine treatment duration must have been given for ≥6 months prior to disease progression. This may be the endocrine treatment component of the CDK4/6 inhibitor line of therapy.</li></ul></li><li>• Radiological progression during or after the last line of therapy.</li><li>• Measurable disease evaluable per Response Evaluation Criterion in Solid Tumors (RECIST) v.1.1 or non-measurable bone-only disease</li><li>• Eastern Cooperative Oncology Group (ECOG) performance status 0-1</li><li>• Participants should be willing to provide blood and tumor tissue</li></ul>
Critères d'exclusion	<ul style="list-style-type: none"><li>• Participants with advanced, symptomatic visceral spread, that are at risk of life-threatening complications in the short term</li><li>• Prior treatment with:<ul style="list-style-type: none"><li>• ARV-471, fulvestrant, elacestrant, mTOR, PI3K, AKT pathway inhibitors, PARP inhibitor for any setting</li><li>• other investigational agents (including novel endocrine therapy any SERDs, SERCAs, CERANs) for any setting</li></ul></li><li>• prior chemotherapy for advanced/metastatic disease</li><li>• Inadequate liver, kidney and bone marrow function</li><li>• Active brain metastases</li><li>• Participants with significant concomitant illness</li></ul>

