



Essai Clinique

Généré le 22 mai 2025 à partir de

| | |
|-------------------------|---|
| Titre | Étude ouverte de phase III à répartition aléatoire visant à comparer l'implunestrant adjuvant et une endocrinothérapie adjuvante de référence chez des patients ayant déjà reçu une endocrinothérapie adjuvante pendant 2 à 5 ans pour le traitement d'un cancer du sein au stade précoce exprimant des récepteurs d'œstrogènes (RE+), HER2- associé à un risque accru de récurrence. |
| Protocole ID | EMBER-4 |
| ClinicalTrials.gov ID | NCT05514054 |
| Type(s) de cancer | Sein |
| Phase | Phase III |
| Type étude | Clinique |
| Médicament | Implunestrant versus thérapie endocrinienne adjuvante standard |
| Institution | CIUSSS DU NORD-DE-L'ILE-DE-MONTREAL HOPITAL DU SACRE-COEUR-DE-MONTREAL 5400 boul. Gouin Ouest, Montréal, QC, H4J1C5 |
| Ville | |
| Investigateur principal | Dre Isabelle Gingras |
| Coordonnateur | Marie-Anne Capobianco 514-338-2222 poste 3493 |
| Statut | Actif en recrutement |
| Date d'activation | 23-10-2023 |
| But étude | The main purpose of this study is to measure how well implunestrant works compared to standard hormone therapy in participants with early breast cancer that is estrogen receptor positive (ER+) and human epidermal receptor 2 negative (HER2-). Participants must have already taken endocrine therapy for two to five years and must have a higher-than-average risk for their cancer to return. Study participation could last up to 10 years. |
| Critères d'éligibilité | <ul style="list-style-type: none">• Have a diagnosis of ER+, HER2- early-stage, resected, invasive breast cancer without evidence of distant metastasis.• Participants must have received at least 24 months but not more than 60 months of any adjuvant ET, from time of adjuvant ET initiation.• Participants may have received (neo) adjuvant chemotherapy and/or targeted therapy with a CDK4/6- or PARP- inhibitor.• Must have an increased risk of disease recurrence based on clinical-pathological risk features.• Have a Performance Status of 0 or 1 on the Eastern Cooperative Oncology Group scale.• Have adequate organ function. |
| Critères d'exclusion | <ul style="list-style-type: none">• Have any evidence of metastatic disease (including contralateral ALN) or inflammatory breast cancer at primary breast cancer diagnosis.• Participants with more than a 6-month consecutive gap in therapy during the course of prior adjuvant ET.• Participants who have completed or discontinued prior adjuvant ET >6 months prior to screening.• Participants with a history of previous breast cancer are excluded, with the exception of ipsilateral DCIS treated by locoregional therapy alone ≥5 years ago.• Pregnant, breastfeeding, or expecting to conceive or father children within the projected duration of the trial, starting with the screening visit through 180 days after the last dose of study intervention. |

- Participant has previously received ET of any duration for breast cancer prevention (tamoxifen or AIs) or raloxifene.
- Participants with a history of any other cancer.
- Have serious preexisting medical conditions that, in the judgment of the investigator, would preclude participation in this study.