

Warning

This fact sheet aims to let you know the side effects most commonly seen following the administration of your treatment. Please note that when you take any medication, there is a possibility that it causes rare side effects and it is difficult to predict which patients will be affected.

General information for patient Bevacizumab

Date: June 2006

Revision: October 2016

Medication	Administration
Bevacizumab	Intravenous infusion on day 1 of each cycle of chemotherapy 1 st infusion : 90 minutes 2 nd infusion: 60 minutes * 3 rd infusion and more: 30 minutes * <i>* if prior dose well tolerated</i>
<hr/> <i>Repeat the treatment every ____ weeks (with chemotherapy)</i> <hr/>	

Warning

This fact sheet aims to let you know the side effects most commonly seen following the administration of your treatment. Please note that when you take any medication, there is a possibility that it causes rare side effects and it is difficult to predict which patients will be affected.

General Information associated with your treatment

- › Bevacizumab is used in combination with chemotherapy to treat certain types of cancer. It is a monoclonal antibody which enhances the efficacy of your chemotherapy.
- › A **blood test, a urinary analysis** as well as a **measurement of your blood pressure** will be done prior to every treatment.
- › Bevacizumab administered during pregnancy may harm the embryo. A method of **contraception** is strongly recommended during treatments and for 6 months after your last dose of bevacizumab. Breast-feeding is contra-indicated during the whole duration of your treatment.
- › Consult your doctor or pharmacist before you take other medications (even the over the counter medications available without a prescription) or natural products. These may provoke an interaction with your treatment and diminish its efficacy or increase side effects.
- › It is very important to notify us if you start taking medications such as Aspirin®, blood thinners (ex: Coumadin®...), anti-inflammatory drugs (ex: Motrin®, Advil®...)
- › Follow the instructions given by your cancer center about the right way to handle and dispose of body fluids (for example urine, stools) in order to protect the environment and your relatives.
- › Be sure to always have at hand the complete list of your medications, dietary supplements, vitamins and other natural health products. Give it to you doctor and your pharmacist. It is important to make them aware of any change.

Warning

This fact sheet aims to let you know the side effects most commonly seen following the administration of your treatment. Please note that when you take any medication, there is a possibility that it causes rare side effects and it is difficult to predict which patients will be affected.

Adverse effects and their management

Adverse effects	Management
Bevacizumab is administered in combination with chemotherapy. We suggest that you consult the pamphlet which describes the specific information regarding your chemotherapy regimen.	
Minor bleeding such as nose bleed may occur.	<ul style="list-style-type: none">› Sit down straight with you head slightly bent forward.› Apply a continuous pressure on your nostrils for a few minutes then check if the bleeding has stopped.› Avoid blowing your nose for 12 hours after the bleeding.› If the bleeding lasts more than 20 minutes, consult immediately your physician or go to the Emergency Room.
High blood pressure might develop.	<ul style="list-style-type: none">› Before each treatment, your nurse will measure your blood pressure.› If you experience high blood pressure, medications can be prescribed to control it.
Abdominal pain may occur.	<ul style="list-style-type: none">› Notify your physician or pharmacist if this happens.› If you experience severe, sudden pain with nausea, vomiting or constipation, go to the Emergency Room immediately.
If you experience any of the following symptoms: Severe headache, visual problems, difficulty talking, loss of coordination, weakness or numbness, chest pain or chest tightness.	<ul style="list-style-type: none">› Go immediately to the Emergency Room.

Special Consideration

- › Notify your dentist or physician that you are receiving bevacizumab if you have to undergo a surgery while on treatment.

Pharmacist: _____ **Phone:** _____
from ____ h to ____ h (Monday to Friday)